CDPH LTC Toolkit 2018

Example 4.10 O'Connor Hospital Pharmacy Communication Sheet for Vancomycin

	Addressograph
O'Connor	
Hospital	
It's The Way We Care	
Member of Daughters of Charity Health System	
ANTIBIOTIC STEWARDSHIP	
PHARMACY COMMUNICATION ST Vancomycin	HEET
TO PHYSICIAN:	DATE:
REGARDING PATIENT	
Please consider discontinuing	g vancomycin in this patient.
CDC recommenda	
Situations in which the use of	vancomycin is discouraged: I infections in patients whose cultures are negative for beta-lactam—
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-asso	vancomycin is discouraged: I infections in patients whose cultures are negative for beta-lactam—
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-asso	vancomycin is discouraged: d infections in patients whose cultures are negative for beta-lactam— s. ciated colitis. Vancomycin should be reserved for when antibiotic—
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-asso	vancomycin is discouraged: d infections in patients whose cultures are negative for beta-lactam— s. ciated colitis. Vancomycin should be reserved for when antibiotic- tronidazole therapy or is severe and potentially life-threatening.
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-asso	vancomycin is discouraged: It infections in patients whose cultures are negative for beta-lactames. Sciated colitis. Vancomycin should be reserved for when antibiotic-tronidazole therapy or is severe and potentially life-threatening. Respectfully,
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-associated colitis fails to respond to metal	vancomycin is discouraged: It infections in patients whose cultures are negative for beta-lactames. Sciated colitis. Vancomycin should be reserved for when antibiotic-tronidazole therapy or is severe and potentially life-threatening. Respectfully,
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-associated colitis fails to respond to metal	vancomycin is discouraged: It infections in patients whose cultures are negative for beta-lactames. Sciated colitis. Vancomycin should be reserved for when antibiotic-tronidazole therapy or is severe and potentially life-threatening. Respectfully,
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-associated colitis fails to respond to metal	vancomycin is discouraged: It infections in patients whose cultures are negative for beta-lactames. Sciated colitis. Vancomycin should be reserved for when antibiotic-tronidazole therapy or is severe and potentially life-threatening. Respectfully,
Situations in which the use of Continued empiric use for presumed resistant Gram-positive microorganisms Eradication of MRSA colonization. Primary treatment of antibiotic-associated colitis fails to respond to metal	vancomycin is discouraged: It infections in patients whose cultures are negative for beta-lactames. Sciated colitis. Vancomycin should be reserved for when antibiotic-tronidazole therapy or is severe and potentially life-threatening. Respectfully,

For more information about this example contact Olga DeTorres PharmD at OlgaDeTorres@verity.org