California Influenza and Other Respiratory Disease Surveillance for Week 43 (October 21–27, 2012)

Note: This report includes data from many different sources of influenza surveillance, including syndromic surveillance, laboratory surveillance, and mandatory reporting of influenza deaths for cases ages 0-64 years. The information contained in this report should be viewed as a "snapshot" of influenza activity for each surveillance week, and should not be considered as population-based data or representative of all California public health jurisdictions.

Based on low levels of influenza-like illness and laboratory detections, overall influenza activity in California during Week 43 was "sporadic*."

Influenza Report Highlights

- Outpatient influenza-like illness (ILI) decreased 1.6% in Week 43 (0.9%) compared to Week 42 (2.5%).
- Of 501 specimens tested during Week 43,
 - 8 (1.6%) were positive for influenza; of these
 - 2 (25.0%) were influenza B and
 - 6 (75.0%) were influenza A.
 - The influenza A specimens were not subtyped.
- The California Department of Public Health Viral and Rickettsial Disease Laboratory (CDPH-VRDL) has not tested any specimens for antiviral resistance to date.
- No California specimens have been strain-typed by the Centers for Disease Control and Prevention (CDC) to date.
- CDPH has received no reports of laboratory-confirmed influenza-associated deaths among persons less than 65 years of age.
- No cases of novel influenza have been detected in California.


*For the Centers for Disease Control and Prevention (CDC) definitions of influenza geographic distribution, please go to the CDC Influenza page (http://www.cdc.gov/flu/weekly/overview.htm).

A. Syndromic Surveillance Update

1.CDC Influenza Sentinel Providers

A total of 66 enrolled sentinel providers have reported data for Week 43, compared to an average of 101 providers reporting for each of the previous weeks. Based on available data, the percentage of ILI visits for Week 43 (0.9%) was below baseline (Figure 1).

Figure 1. Percentage of Influenza-like Illness Visits Among Patients Seen by California Sentinel Providers, 2008–2012




The seasonal baseline was calculated using a regression model applied to data from the previous seven years. The epidemic threshold is two standard deviations above the seasonal baseline and is the point at which the observed percentage of ILI is significantly higher than would be expected at that time of the year.

Kaiser Permanente Hospitalization Data

The percentage of hospitalizations for pneumonia and influenza (P&I) in Kaiser Permanente facilities in northern California was below baseline during Week 43 (Figure 2). During Week 43, 3.5% of hospitalizations were due to P&I, compared to 3.7% of hospitalizations in the previous week.

Figure 2. Percentage of P&I Hospitalizations in Kaiser Permanente Northern California Hospitals, 2008–2012



The seasonal baseline was calculated using a regression model applied to data from the previous five years. The epidemic threshold is two standard deviations above the seasonal baseline and is the point at which the observed percentage of pneumonia and influenza hospitalizations in Kaiser Permanente hospitals in northern California is significantly higher than would be expected at that time of the year.

B. Laboratory Update

1. Respiratory Laboratory Network (RLN) and Sentinel Laboratory Surveillance Results

The percentage of influenza detections in the RLN and sentinel laboratories remained low in Week 43 (1.6%) (Figure 3). In Week 43, of 501 specimens tested by the RLN and sentinel laboratories, 2 (0.4%) were positive for influenza B and 6 (1.2%) were positive for influenza A. The specimens that tested positive for influenza A had no further subtyping performed. The influenza-positive specimens were reported from Alameda, Los Angeles, San Francisco, and Santa Clara counties.

To date for the 2012-2013 season, of 1,864 specimens tested, 19 (1.0%) were positive for influenza; of these, 6 (31.6%) were influenza B and 13 (68.4%) were influenza A. Of the 13 specimens that tested positive for influenza A, 4 (30.8%) were subtyped as seasonal A (H3) and 9 (69.2%) had no further subtyping performed.


Neither the RLN nor CDPH-VRDL have identified any influenza viruses by polymerase chain reaction (PCR) typing or subtyping that are suggestive of a novel influenza virus.

60% 2007-2008 -- 2008-2009 --2009-2010 55% Percentage of respiratory specimens in which 50% 2010-2011 2011-2012 2012-2013 45% influenza virus was identified 40% 35% 30% 25% 20% 15% 10% 5% 0% 40 42 44 46 48 50 52 2 6 8 20 22 24 26 28 30 32 34 36 38 10 12 14 16 18 Oct Nov Ďес Jan Feb Mar Apr May Jun Jul Sept Aug Week

Figure 3. Percentage of Influenza Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2007–2012

The proportion of respiratory syncytial virus (RSV) detections increased in Week 43 (3.9%, compared to 1.9% in Week 42) (Figure 4).

Figure 4. Percentage of RSV Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2007–2012



Antiviral Resistance Testing

CDPH-VRDL has not tested any influenza specimens for antiviral resistance to date during the 2012-2013 influenza season.

2. Influenza Virus Strain Characterization

No California specimens have been strain-typed to date during the 2012-2013 influenza season.

C. Laboratory-confirmed Fatal Case Reports

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among cases age 0- 64 years with laboratory-confirmed influenza are reportable to CDPH.

CDPH has received no reports of laboratory-confirmed influenza-associated deaths among patients under 65 years of age during the 2012–2013 influenza season.

D. Influenza-associated Outbreaks

CDPH has received no reports of laboratory-confirmed influenza outbreaks during the 2012–2013 influenza season.

For questions regarding influenza surveillance and reporting in California, please email lnfluenzaSurveillance@cdph.ca.gov. This account is monitored daily by several epidemiologists.

For more information regarding the different influenza surveillance data sources, please visit the CDPH Influenza Surveillance Program at

https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Flu-Reports.aspx

To obtain additional information regarding influenza, please visit the <u>CDPH influenza website</u> at https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx.

A copy of the case report form for reporting any laboratory-confirmed influenza case that was either admitted to the ICU or died can be downloaded from the Severe Influenza Case History Form Link at https://www.cdph.ca.gov/CDPH%20Document%20Library/ControlledForms/cdph9070.pdf.