
Influenza and Other Respiratory Viruses Weekly Report

California Influenza Surveillance Program

Highlights (Week 42: October 13, 2019 – October 19, 2019)

Statewide Activity



Note: This report includes data from many sources of influenza surveillance and it should be viewed as a preliminary "snapshot" of influenza activity for each surveillance week. Because data are preliminary, the information may be updated in later reports as additional data are received. These data should not be considered population-based or representative of all California public health jurisdictions.

A. Outpatient, Inpatient, and Death Data

1. Influenza Sentinel Providers

Sentinel providers (physicians, nurse practitioners, and physician assistants) situated throughout California report on a weekly basis the number of patients seen with influenza-like illness (ILI) and the total number of patients seen for any reason. ILI is defined as any illness with fever (≥100°F or 37.8°C) AND cough and/or sore throat (in the absence of a known cause other than influenza).

A total of 87 enrolled sentinel providers have reported data for Week 42. Based on available data, the percentage of visits for ILI during Week 42 was 1.8% compared to Week 41 (1.6%) and is within expected levels for this time of year (Figure 1).

Percent II I - - Baseline II I Two Standard Deviations above Baseline 9 Percentage of Influenza-Like Illness Visits 8 7 6 5 4 3 101012017 21,012018 410/2018 Grozoro 811012016 10/10/2016 12/10/2016 2110/2017 MOROTT Ghoron1 8/10/2017 121102017 6/10/2018 8/10/2018 10102018 1211012018 21/0/2019 410/2019 611012019 811012019 10/10/2019 1211012019 Week Ending Date

Figure 1. Percentage of Influenza-like Illness Visits Among Patients Seen by California Sentinel Providers, 2015–2020


The seasonal baseline was calculated using a regression model applied to data from the previous five years. Two standard deviations above the seasonal baseline is the point at which the observed percentage of ILI is significantly higher than would be expected at that time of year.

2. Kaiser Permanente Hospitalization Data

Inpatients at Kaiser Permanente facilities with an admission diagnosis including the keywords "flu," "influenza," "pneumonia," or variants of the keywords are defined as pneumonia and influenza (P&I)-related admissions. The number of P&I admissions is divided by the total number of hospital admissions occurring in the same time period to estimate the percentage of P&I admissions. Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are excluded from the denominator.

The percentage of admissions for pneumonia and influenza (P&I) in Kaiser Permanente facilities in northern California during Week 42 was 3.6% compared to Week 41 (3.9%) and is within expected levels for this time of the year (Figure 2).

Figure 2. Percentage of P&I Admissions in Kaiser Permanente Northern California Hospitals, 2015–2020




The seasonal baseline was calculated using a regression model applied to data from the previous five years. Two standard deviations above the seasonal baseline and is the point at which the observed percentage of pneumonia and influenza hospitalizations in Kaiser Permanente hospitals in Northern California is significantly higher than would be expected at that time of the year.


The majority of admissions for pneumonia and influenza did not result in intensive care unit (ICU) admission or death; however, 42 ICU admissions and 25 deaths have occurred among persons with P&I admission diagnoses (Figure 3a). The majority of P&I admissions occurred among persons ≥65 years of age across all severity categories, especially among deaths (Figure 3b). Please note that pneumonia and influenza admissions serve as a proxy for influenza activity, but do not necessarily represent influenza infections.

Figure 3. Number (a) and age group distribution (b) of non-ICU, ICU, and Deaths associated with P&I Admissions in Kaiser Permanente Northern California Hospitals, 2019–2020 Season to Date

(a)



(b)




4. Influenza Mortality Surveillance from Death Certificates

Deaths occurring in California among residents who had influenza noted in any cause of death field on the death certificate (text or coded) are defined as influenza-coded deaths. The percentage of influenza-coded deaths is calculated by dividing the number of influenza-coded deaths by the total number of all cause deaths during the same period. Influenza-coded deaths are not necessarily laboratory-confirmed and are an underestimate of all influenza-associated deaths.


During Week 42, no new influenza-coded deaths were identified. To date during the 2019–2020 influenza season, one influenza-coded death has been identified (Figure 4). The percentage of deaths coded as influenza during Week 42 was 0.0% compared to 0.0% during Week 41 (Figure 5).

Figure 4. Number of Influenza-coded Deaths Identified from Death Certificates by Week of Death, 2019–2020 Season



Note: Coding of deaths can be delayed by several weeks. Influenza-coded deaths will be included once enough information is available to identify them.

Figure 5. Percentage of Influenza-coded Deaths Occurring in California among California Residents, 2015–2020



5. Laboratory-Confirmed Severe Influenza-associated Pediatric Deaths

Influenza-associated deaths in children <18 years of age are nationally notifiable. The weekly influenza report includes confirmed deaths formally reported to CDPH through October 19, 2019 (Week 42).

No laboratory-confirmed influenza-associated fatalities in children <18 years of age were reported to CDPH during Week 42. To date, CDPH has received no reports of laboratory-confirmed influenza-associated deaths among persons <18 years of age during the 2019–2020 influenza season.

B. Laboratory Update – Influenza

1. Respiratory Laboratory Network (RLN) and Clinical Sentinel Laboratory Surveillance Results

Laboratory surveillance for influenza and other respiratory viruses involves the use of data from clinical sentinel laboratories (hospital, academic, and private laboratories) and public health laboratories in the Respiratory Laboratory Network located throughout California. These laboratories report the number of laboratory-confirmed influenza and other respiratory virus detections and isolations on a weekly basis.

The overall percentage of influenza detections in clinical sentinel laboratories in Week 42 (2.1%) was higher than Week 41 (1.8%) (Figure 6). Additional details, including influenza typing and subtyping information from public health laboratories can be found in Figures 6 and 7 and Tables 1 and 2.

Neither the RLN nor CDPH-VRDL has identified any influenza viruses by polymerase chain reaction (PCR) that are suggestive of a novel influenza virus.

Figure 6. Percentage of Influenza Detections at Clinical Sentinel Laboratories, 2015–2020




Figure 7. Number of Influenza Detections by Type and Subtype Detected in the Respiratory Laboratory Network, 2019–2020




Table 1. Respiratory Specimens Testing Positive for Influenza — Clinical Sentinel Laboratories, Current Week and Season to Date

	Current Week Number	Current Week Percent	Season to Date Number	Season to Date Percent
Number of Specimens Tested	1,951		6,141	
Influenza Positive	40	2.1	108	1.8
Α	26	65.0*	71	65.7 [*]
В	14	35.0*	37	34.3*

^{*} Percent of specimens positive for influenza

Table 2. Respiratory Specimens Testing Positive for Influenza by Influenza Type and Subtype — Respiratory Laboratory Network, Current Week and Season to Date

	Current Week Number	Current Week Percent	Season to Date Number	Season to Date Percent
Influenza Positive	9		46	
Α	8	88.9*	34	73.9 [*]
A (H1)pdm09	2	25.0 [†]	9	26.5 [†]
A (H3)	6	75.0 [†]	24	70.6 [†]
A, not subtyped	0	0.0†	1	2.9 [†]
В	1	11.1*	12	26.1 [*]
B Victoria	0	0.0‡	11	91.7 [‡]
B Yamagata	0	0.0‡	0	0.0‡
B, not lineage typed	1	100.0‡	1	8.3 [‡]

^{*} Percent of specimens positive for influenza


D. Influenza-Associated Outbreaks

No laboratory-confirmed influenza outbreaks were reported to CDPH during Week 42. To date, no laboratory-confirmed influenza outbreaks have been reported to CDPH for the 2019–2020 season.

[†] Percent of influenza A positives

[‡] Percent of influenza B positives

Figure 8. Number of Laboratory-Confirmed Influenza-Associated Outbreaks by Week of First Onset, 2018–2020




 * Earliest date associated with the outbreak was used for outbreaks without reported date of first patient's symptom onset.


E. California Border Region Influenza Surveillance Network Data

The border influenza surveillance network is comprised of outpatient provider sentinel sites whose geographical coverage extends approximately 100 kilometers (60 miles) north of the California-Baja California border and includes Imperial and San Diego Counties, as well as some parts of Riverside County.

1. Syndromic Surveillance Update

A total of 9 border region sentinel providers reported data during Week 42. The total number of patients screened by all sentinel sites for ILI during Week 42 was 7,519. Outpatient ILI activity was 0.3% in Week 42. ILI activity for the California border region during Week 42 was lower when compared to activity for the same week during the 2017–2018 season and 2018–2019 season (Figure 9). All influenza syndromic data summarized for the border region represent a subset of CDC influenza sentinel providers in California.






2. Virologic Surveillance Update

During Week 42, 160 respiratory specimens were tested from border region sentinel clinical laboratories; of these, one (0.6%) tested positive for influenza (1 [100.0%] influenza A). Cumulatively this season, a total of 593 respiratory specimens were tested from border region sentinel clinical laboratories; of these, six (1.0%) tested positive for influenza (5 [83.3%] influenza A; 1 [16.7%] influenza B).

During Week 42, 0 influenza positive specimens were detected at border region RLN laboratories. Cumulatively this season, a total of four influenza positive specimens have been detected at border region RLN laboratories; of which, four (100.0%) were influenza A. Of the four specimens that tested positive for influenza A at RLN laboratories, two (50.0%) were subtyped as A (H1)pdm09 and two (50.0%) were subtyped as A (H3).

Laboratory data summarized in Figure 10 include data from border region influenza clinical sentinel laboratories (percentage of specimens testing positive for influenza) as well as data from border region RLN laboratories (influenza type and subtype/lineage type).

Figure 10. Number of Influenza Detections by Type and Subtype Detected in Respiratory Laboratory Network Laboratories and the Percentage of Specimens Testing Positive at Clinical Sentinel Laboratories — California Border Region, 2019–2020



F. Other Respiratory Viruses

1. Laboratory-Confirmed Severe Respiratory Syncytial Virus Case Reports

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among children aged 0–4 years with laboratory-confirmed respiratory syncytial virus (RSV) are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH through October 19, 2019 (Week 42).

No laboratory-confirmed RSV-associated deaths among children <5 years of age were reported to CDPH during Week 42. To date, CDPH has received no reports of laboratory-confirmed RSV-associated deaths among children <5 years of age during the 2019–2020 influenza season.

2. Other Respiratory Virus Laboratory Update

During Week 42, 1,877 specimens were tested for RSV and 13 (0.7%) were positive, which is lower than Week 41 (1.1%) (Figure 11). During Week 42, coronavirus, parainfluenza, rhinovirus/enterovirus, and human metapneumovirus virus activity increased; and adenovirus activity decreased. (Figure 12).

Figure 11. Percentage of RSV Detections at Clinical Sentinel Laboratories, 2015–2020





Figure 12. Percentage of Other Respiratory Pathogen Detections at Clinical Sentinel Laboratories, 2019–2020



Activity Levels:

No Activity: No laboratory-confirmed cases of influenza and no reported increase in the number of cases of ILI.

Sporadic: Small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of ILI.

Local: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in a single region of the state.

Regional: Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions.

Widespread: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

California Regions:

Northern: Alpine, Amador, Butte, Colusa, Del Norte, El Dorado, Glenn, Humboldt, Lake, Lassen, Mendocino, Modoc, Nevada, Placer, Plumas, Sacramento, Shasta, Sierra, Siskiyou, Sutter, Tehama, Trinity, Yolo, and Yuba counties

Bay Area: Alameda, Contra Costa, Marin, Napa, Solano, San Francisco, San Mateo, Santa Clara, Santa Cruz, and Sonoma counties

Central Valley: Calaveras, Fresno, Inyo, Kings, Mono, Madera, Mariposa, Merced, Monterey, San Benito, San Joaquin, Stanislaus, Tulare, and Tuolumne counties **Upper Southern:** Kern, Los Angeles, San Luis Obispo, Santa Barbara, and Ventura counties

Lower Southern: Imperial, Orange, Riverside, San Bernardino, and San Diego counties

For questions regarding influenza surveillance and reporting in California, please email lnfluenzaSurveillance@cdph.ca.gov. This account is monitored daily by several epidemiologists.

To obtain additional information regarding influenza, please visit the <u>CDPH influenza</u> website

(https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx).

A copy of the case report form for reporting any laboratory-confirmed influenza case that was either admitted to the ICU or died can be downloaded from the <u>CDPH influenza</u> website

(https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx).

For information about national influenza activity, please visit the Centers for Disease Control and Prevention's <u>FluVlew</u> (https://www.cdc.gov/flu/weekly/index.htm) and <u>FluView Interactive</u> (https://www.cdc.gov/flu/weekly/fluviewinteractive.htm) websites.