




ZIKA SCREENING ALGORITHM



FOR CHILDREN AND ADULTS



FOR INFANTS

INFANT ZIKA VIRUS TESTING FOR SUSPECTED CONGENITAL ZIKA VIRUS INFECTION SHOULD BE DONE IN THE FOLLOWING CASES:

- Maternal laboratory evidence of Zika virus infection
- Infant findings consistent with Congenital Zika Virus Syndrome and maternal exposure (such as travel or sexual contact) regardless of maternal test results
- Maternal exposure, but mother not tested or results were inconclusive or unavailable before delivery, even if infant appears healthy; in such cases, newborn testing may be done in parallel with maternal testing

See CDC documents for newborn management and first year recommendations at: www.cdc.gov/zika/pdfs/zika_peds.pdf and www.cdc.gov/zika/pdfs/pediatric-evaluation-follow-up-tool.pdf

See CDPH Evaluation and Follow-Up Procedures for Suspected Congenital Zika Virus Infection at: www.cdph.ca.gov/HealthInfo/discond/Documents/GuidanceforCongenitalZikaInfectionEvaluation.pdf

See CDPH guidance for lab testing: www.cdph.ca.gov/programs/vrdl/Documents/Zika_Testing_VRDL_Quicksheet.pdf

For questions about Zika virus testing or test results, contact your local health department: www.cdph.ca.gov/programs/cclho/Documents/LHD_CD_Contact_Info_1-28-16.pdf