

California Department of Public Health Influenza Surveillance Program

California Influenza and Other Respiratory Disease Surveillance for Week 48

(November 27, 2016 to December 3, 2016)

Note: This report includes data from many sources of influenza surveillance and it should be viewed as a preliminary “snapshot” of influenza activity for each surveillance week. Because data are preliminary, the information may be updated in later reports as additional data are received. These data should not be considered population-based or representative of all California public health jurisdictions.

Overall influenza activity in California remained “sporadic*” during Week 48.

Influenza Report Highlights

- Influenza activity in California is within expected activity levels for this time of year
- Outpatient influenza-like illness (ILI)
 - 1.1% of patient visits during Week 48 were for ILI, which is within expected levels for this time of year
- Hospitalization data
 - 4.6% of Kaiser patients hospitalized during Week 48 were admitted with a pneumonia or influenza (P&I) diagnosis, which is within expected levels for this time of year
- Influenza virus detections by Respiratory Laboratory Network and Sentinel Laboratories
 - 150 (6.3%) of 2,378 specimens tested were positive for influenza during Week 48
- Influenza-associated deaths among patients 0–64 years of age
 - One laboratory-confirmed influenza death was reported during Week 48
- Influenza-associated outbreaks
 - One laboratory-confirmed influenza outbreak was reported during Week 48

***Sporadic:** Small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of ILI.


A. Outpatient and Inpatient Data

1. Influenza Sentinel Providers

Sentinel providers (physicians, nurse practitioners, and physician assistants) situated throughout California report on a weekly basis the number of patients seen with influenza-like illness (ILI) and the total number of patients seen for any reason. ILI is defined as any illness with fever ($\geq 100^{\circ}\text{F}$ or 37.8°C) AND cough and/or sore throat (in the absence of a known cause other than influenza).

A total of 83 enrolled sentinel providers have reported data for Week 48. Based on available data, the percentage of visits for ILI during Week 48 was 1.1% compared to Week 47 (2.2%) and was within expected baseline levels for this time of year (Figure 1).

Figure 1. Percentage of Influenza-like Illness Visits Among Patients Seen by California Sentinel Providers, 2012–2017




2. Kaiser Permanente Hospitalization Data

Inpatients at Kaiser Permanente facilities with an admission diagnosis including the keywords “flu,” “influenza,” “pneumonia,” or variants of the keywords are defined as pneumonia and influenza (P&I) related admissions. The number of P&I admissions is divided by the total number of hospital admissions occurring in the same time period to estimate the percentage of P&I admissions. Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are excluded from the denominator.

The percentage of hospitalizations for pneumonia and influenza (P&I) in Kaiser Permanente facilities in northern California during Week 48 was 4.6% compared to Week 47 (4.1%) and was within expected levels for this time of the year (Figure 2).

Figure 2. Percentage of P&I Hospitalizations in Kaiser Permanente Northern California Hospitals, 2012–2017




3. Influenza-Associated Hospitalizations, California Emerging Infections Program

The California Emerging Infections Program (CEIP), Influenza Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-associated hospitalizations among patients of all ages in Alameda, Contra Costa, and San Francisco counties.

The incidence of influenza-associated hospitalizations per 100,000 population decreased in Week 47 (0.17) compared to Week 46 (0.28) (Figure 3). Data for Week 48 are not presented because results are still being collected and are likely to change.

Figure 3. Incidence of Influenza Hospitalizations in CEIP Counties, 2014–2017



Note: The 2014-15 season contains a week 53. Data have been shifted so that week 1 aligns across years.

B. Laboratory Update - Influenza


1. Respiratory Laboratory Network (RLN) and Sentinel Laboratory Surveillance Results

Laboratory surveillance for influenza and other respiratory viruses involves the use of data from hospital, academic, private and public health laboratories located throughout California. These laboratories report the number of laboratory-confirmed influenza and other respiratory virus detections and isolations on a weekly basis.

The percentage of influenza detections in the RLN and sentinel laboratories in Week 48 (6.3%) was higher than Week 47 (4.9%) (Figure 3). Additional details can be found in Figures 4 and 5 and Table 1.

Neither the RLN nor CDPH-VRDL have identified any influenza viruses by polymerase chain reaction (PCR) that are suggestive of a novel influenza virus.

Figure 4. Percentage of Influenza Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2012–2017



Note: The 2014–15 season contains a week 53. Data have been shifted so that week 1 aligns across years.

Figure 5. Number of Influenza Detections by Type and Subtype Detected in Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017




Table 1. Respiratory Specimens Testing Positive for Influenza by Influenza Type and Subtype - Respiratory Laboratory Network and Sentinel Laboratories, Current Week and Season to Date

	Week 48 (Number)	Week 48 (Percent)	Season to Date (Number)	Season to Date (Percent)
Number of Specimens Tested	2,378		18,800	
Number of Specimens Positive for Influenza	150	6.3*	720	3.8*
Influenza Type/Subtype of Positive Specimens				
A	142	94.7 [†]	685	95.1 [†]
2009 A (H1)	3	2.1 [‡]	7	1.0 [‡]
A (H3)	39	27.5 [‡]	171	27.5 [‡]
A, not subtyped	100	70.4 [‡]	507	74.0 [‡]
B	8	5.3 [†]	35	4.9 [†]

* Percent of total specimens tested for influenza

[†] Percent of specimens positive for influenza

[‡] Percent of influenza A positives

2. Antiviral Resistance Testing

Of the influenza specimens tested by the CDPH-VRDL to date this season, no specimens have been found to be resistant to Oseltamivir (Table 2).

Table 2. Number of specimens tested for Oseltamivir resistance, 2016–2017

	Oseltamivir Resistance
Influenza 2009A (H1)	0/2
Influenza A (H3)	0/2
Influenza B	0/3

3. Influenza Virus Strain Characterization

No California specimens have been strain-typed to date during the 2016–2017 influenza season.

C. Laboratory-Confirmed Severe Influenza Case Reports

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among patients aged 0–64 years with laboratory-confirmed influenza are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH as of December 3, 2016 (Week 48).

During Week 48, one laboratory-confirmed influenza fatality was reported. To date, CDPH has received one report of a laboratory-confirmed influenza-associated death among patients <65 years of age during the 2016–2017 influenza season.

D. Influenza-Associated Outbreaks

During Week 48, one laboratory-confirmed influenza outbreaks was reported. To date, six laboratory-confirmed influenza outbreaks have been reported to CDPH for the 2016–2017 season.

E. California Border Region Influenza Surveillance Network Data

The border influenza surveillance network is comprised of outpatient provider sentinel sites whose geographical coverage extends approximately 100 kilometers (60 miles) north of the California-Baja California border and includes Imperial and San Diego Counties, as well as some parts of Riverside County.


Syndromic Surveillance Update

A total of 14 border region sentinel providers reported data during Week 48, compared to 15 during Week

47 of 2016. The total number of patients screened by all sentinel sites for ILI during Week 48 was 12,491.

Outpatient ILI activity was lower in Week 48 (0.7%) compared to Week 47 (1.0%). ILI activity for the California border region during Week 48 was lower when compared to activity for the same week during the 2014–2015 and 2015–2016 season (Figure 6). All influenza syndromic data summarized for the border region represents a subset of CDC influenza sentinel providers in California.

Figure 6. Percentage of Influenza-like Illness Visits among Patients Seen by California Border Region Sentinel Providers, 2014–2017




Note: The 2014-15 season contains a week 53. Prior years' data have been shifted so that week 1 aligns across years.

Virologic Surveillance Update

Cumulatively this season, a total of 1,919 respiratory specimens have been tested from border region clinics; of these, 37 (1.9%) tested positive for influenza. Of the 37 specimens that tested positive, 32 (86.5%) were influenza A and 5 (13.5%) were influenza B. Of the 32 specimens that tested positive for influenza A, 3 (9.4%) were subtyped as 2009 A (H1), 19 (59.4%) were subtyped as A (H3), and 10 (31.3%) had no further subtyping performed. None of the specimens that tested positive for influenza B were lineage typed. For Week 48, a total of 249 respiratory specimens were submitted for testing; 8

(3.2%) were positive for influenza. Of the 8 specimens that tested positive for influenza, 7 (87.5%) were influenza A and 1 (12.5%) was influenza B. Of the 7 specimens that tested positive for influenza A, 1 (14.3%) was subtyped as 2009 A (H1), 5 (71.4%) were subtyped as A (H3), and 1 (14.3%) had no further subtyping performed. None of the specimens that tested positive for influenza B were lineage typed. Laboratory data summarized in Figure 7 includes data from influenza sentinel sites as well as laboratory data from other border region laboratories.

Figure 7. Number of Influenza Detections by Type and Subtype Detected in California Border Region Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017



F. Other Respiratory Viruses

1. Laboratory-Confirmed Severe Respiratory Syncytial Virus Case Reports

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among children aged 0–4 years with laboratory-confirmed respiratory syncytial virus (RSV) are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH as of December 3, 2016 (Week 48).

During Week 48, no laboratory-confirmed RSV fatalities were reported. To date, CDPH has received no reports of laboratory-confirmed RSV-associated deaths among children <5 years of age during the 2016–2017 influenza season.

2. Laboratory Update

During Week 48, 1,850 specimens were tested for RSV and 201 (10.9%) were positive, which was higher than Week 46 (7.1%) (Figure 8). Information on other respiratory viruses can be found in Figure 9.

Figure 8. Percentage of RSV Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2012–2017





Figure 9. Percentage of Other Respiratory Pathogen Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017



For questions regarding influenza surveillance and reporting in California, please email InfluenzaSurveillance@cdph.ca.gov. This account is monitored daily by several epidemiologists.

For more information regarding the different influenza surveillance data sources, please visit the [CDPH Influenza Surveillance Program](#).

To obtain additional information regarding influenza, please visit the [CDPH Influenza Website](#).

Download a copy of the [case report form](#) for reporting any laboratory-confirmed influenza case that was either admitted to the ICU or died.