

## Healthcare-Associated Infections Program Adherence Monitoring **Enhanced Standard Precautions (ESP)**

Assessment completed by:							
Date:							
Unit:							

Regular monitoring with feedback of results to staff can maintain or improve adherence to ESP practices. Use this tool to identify gaps and opportunities for improvement. Monitoring may be performed in any type of resident care location where residents are on ESP. Refer to the grid at the back of this tool to determine if ESP is indicated for the resident.

**Instructions:** Observe 3-4 residents on ESP. Observe each practice and check a box if adherent, Yes or No. In the column on the right, record the total number of "Yes" for adherent practices observed and the total number of observations ("Yes" + "No"). Calculate adherence percentage in the last row.

	Fuhancad Standard Ducas stiens Ducations		ESP		ESP		ESP		ESP		Adherence by Task	
Enhanced Standard Precautions Practices		Resident 1		Resident 2		Resident 3		Resident 4		# Yes	# Observed	
ESP1.	PPE including gloves and gowns are available and located near point of use.	Yes	□No	Yes	□No	Yes	□No	□Yes	□No			
ESP2.	6 Moment/ESP sign is clear and visible. Sample signage is available at <u>CDPH website</u> : (cdph.ca.gov/Programs/CHCQ/HAI/Pages/SNF_EstablishingIC_Program.aspx)	Yes	□No	∐Yes	□No	Yes	□No	Yes	□No			
ESP3.	If a private room is not available, resident is cohorted with a compatible roommate (such as a resident with the same MDRO or resistance mechanism, if present and known).	Yes	□No	Yes	□No	∐Yes	□No	☐Yes	□No			
ESP4.	If multi-bed room, each resident's bed space is treated as a separate room (e.g., care provider performs hand hygiene and changes PPE between caring for each resident in the room).	Yes	□No	Yes	□No	∐Yes	□No	Yes	□No			
ESP5.	Hand hygiene is performed before entering the resident care environment.	☐Yes	□No	☐Yes	□No	Yes	□No	□Yes	□No			
ESP6.	Gloves, gown and other necessary PPE are donned before performing high-contact tasks (6 moments).	Yes	□No	Yes	□No	Yes	□No	□Yes	□No			
ESP7.	Hand hygiene is performed after PPE is removed, before leaving the resident care environment.	Yes	□No	Yes	□No	Yes	□No	☐Yes	□No			
ESP8.	Dedicated or disposable noncritical patient-care equipment (e.g. blood pressure cuffs) is used; if dedicated/disposable equipment is unavailable, then equipment is cleaned and disinfected prior to use on another resident according to manufacturers' instructions.	∏Yes	□No	∏Yes	□No	∏Yes	□No	∐Yes	□No			
# of Correct Practices Observed ("# Yes"):   Total # ESP Observations ("# Observed"):   (Up to 32 total)  (Total "# Yes" ÷ Total "#  If practice could not be observed (i.e. cell is blank), do not count in total # Observed.										% ved" x 100)		

## Enhanced Standard Precautions, including appropriate PPE, should be used for all residents with the following characteristics:



**Indwelling Devices Present (including Ventilator)** 

**Unhealed Wound** 

**Habitual (Persistent or chronic) Incontinence** 

**Complete Dependence (ADLs)** 





- 1. Morning and Evening Care
- 2. Toileting and changing incontinence briefs
- 3. Caring for devices & giving medical treatments
- 4. Wound care
- 5. Mobility assistance & preparing to leave the room
- 6. Cleaning the environment



